

On the strict comparison theorem for G -expectations

Xinpeng LI

School of Mathematics, Shandong University
250100, Jinan, China

January 20, 2010

Abstract. This paper investigates the strict comparison theorem under the framework of G -expectation, i.e., let $X \leq Y$ q.s., if X, Y satisfy some additional conditions, then $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$.

1 Introduction

In 2006, to deal with the model uncertainty problem in finance, Peng establish the sublinear expectation theory and introduce a new sublinear expectation, called G -expectation, which have many well properties as classical linear expectation except linearity (see Peng [10, 11, 12, 13, 14]). Unlike the well-known g -expectation, G -expectation was introduced via fully nonlinear parabolic partial differential equations.

In this paper, we consider the following problem: for a linear expectation E_P , we know that if $X \leq Y$ then $E_P[X] \leq E_P[Y]$, furthermore, if $X \leq Y$ and $P(X < Y) > 0$, then $E_P[X] < E_P[Y]$. If we replace the linear expectation E_P by Peng's G -expectation $\hat{\mathbb{E}}$, the former holds obviously, we interest that when the latter holds. We give three forms of strict comparison theorem for G -expectation and some interesting examples.

This paper is organized as follows: in Section 2, we recall some basic notions and results of G -expectation. In Section 3, we prove strict comparison theorems for G -expectation and give some interesting examples.

2 Preliminaries

We present some preliminaries in the theory of G -expectations. More details can be found in Peng [10, 11, 12, 13, 14].

Let Ω be a given set and let $\mathcal{H}$ be a linear space of real valued functions defined on Ω satisfying: if $X_i \in \mathcal{H}$, $i = 1, \dots, d$, then

$$\varphi(X_1, \dots, X_d) \in \mathcal{H}, \quad \forall \varphi \in C_{l,Lip}(\mathbb{R}^d),$$

where $C_{l,Lip}(\mathbb{R}^d)$ is the space of all real continuous functions defined on $\mathbb{R}$ such that

$$|\varphi(x) - \varphi(y)| \leq C(1 + |x|^k + |y|^k)|x - y|, \forall x, y \in \mathbb{R}^d, \quad k \text{ depends on } \varphi.$$

A sublinear expectation $\mathbb{E}$ on $\mathcal{H}$ is a functional $\mathbb{E} : \mathcal{H} \mapsto \mathbb{R}$ satisfying the following properties:

- (1) Monotonicity: If $X \geq Y$ then $\mathbb{E}[X] \geq \mathbb{E}[Y]$.
- (2) Constant preserving: $\mathbb{E}[c] = c$.
- (3) Sub-additivity: $\mathbb{E}[X + Y] \leq \mathbb{E}[X] + \mathbb{E}[Y]$.
- (4) Positive homogeneity: $\mathbb{E}[\lambda X] = \lambda \mathbb{E}[X], \quad \forall \lambda \geq 0$.

Definition 2.1 In a sublinear expectation space $(\Omega, \mathcal{H}, \mathbb{E})$, a random variable Y is said to be independent from another random variable X if

$$\mathbb{E}[\varphi(X, Y)] = \mathbb{E}[\mathbb{E}[\varphi(x, Y)]_{x=X}], \quad \forall \varphi \in C_{l,Lip}(\mathbb{R} \times \mathbb{R}).$$

Two random variables X_1 and X_2 are called identically distributed, denoted by $X_1 \sim X_2$, if

$$\mathbb{E}[\varphi(X_1)] = \mathbb{E}[\varphi(X_2)], \quad \forall \varphi \in C_{l,Lip}(\mathbb{R}).$$

If $\bar{X}$ is identically distributed with X and independent from X , then $\bar{X}$ is said to be an independent copy of X .

Definition 2.2 In a sublinear expectation space $(\Omega, \mathcal{H}, \mathbb{E})$, a random variable $X \in \mathcal{H}$ is said to be G -normal distributed with $\bar{\sigma}^2 = \mathbb{E}[X^2]$ and $\underline{\sigma}^2 = -\mathbb{E}[-X^2]$, denoted by $X \sim \mathcal{N}(0, [\underline{\sigma}^2, \bar{\sigma}^2])$, if

$$aX + b\bar{X} \sim \sqrt{a^2 + b^2}X, \quad \forall a, b \geq 0,$$

where $\bar{X}$ is an independent copy of X .

We can give a characterization of G -normal distribution by fully nonlinear parabolic partial differential equation (see Peng [10, 11, 12, 13, 14]).

Proposition 2.3 Let $X \sim \mathcal{N}(0, [\underline{\sigma}^2, \bar{\sigma}^2])$, then for each $\varphi \in C_{l,Lip}(\mathbb{R})$, the function u defined by

$$u(t, x) := \mathbb{E}[\varphi(x + \sqrt{t}X)], \quad t \geq 0, x \in \mathbb{R},$$

is the unique viscosity solution of the following G -heat equation:

$$\begin{cases} \partial_t u - G(\partial_{xx} u) = 0, & (t, x) \in [0, \infty) \times \mathbb{R}, \\ u|_{t=0} = \varphi, \end{cases} \quad (1)$$

where $G(\alpha) = (\bar{\sigma}^2 \alpha^+ - \underline{\sigma}^2 \alpha^-)/2$.

We will give the notion of G -expectations. Let $\Omega = C_0(\mathbb{R}^+)$ be the space of all $\mathbb{R}$ -valued continuous paths $(\omega_t)_{t \in \mathbb{R}^+}$ with $\omega_0 = 0$. We consider the canonic process: $B_t(\omega) = \omega_t, t \in [0, \infty), \omega \in \Omega$. For each fixed $T > 0$, we set

$$Lip(\Omega_T) := \{\varphi(B_{t_1}, B_{t_2}, \dots, B_{t_n}) : \forall n \geq 1, t_1, \dots, t_n \in [0, T], \forall \varphi \in C_{l, Lip}(\mathbb{R}^n)\}.$$

and

$$Lip(\Omega) := \bigcup_{n=1}^{\infty} Lip(\Omega_n).$$

Then we can construct a consistent sublinear expectation called G -expectation $\hat{\mathbb{E}}[\cdot]$ on $Lip(\Omega)$, such that B_1 is G -normal distributed under $\hat{\mathbb{E}}$ and for each $s, t \geq 0$ and $t_1, \dots, t_N \in [0, t]$, we have

$$\hat{\mathbb{E}}[\varphi(B_{t_1}, \dots, B_{t_N}, B_{t+s} - B_t)] = \hat{\mathbb{E}}[\psi(B_{t_1}, \dots, B_{t_N})],$$

where $\psi(x_1, \dots, x_N) = \hat{\mathbb{E}}[\varphi(x_1, \dots, x_N, \sqrt{s}B_1)]$. Under G -expectation $\hat{\mathbb{E}}[\cdot]$, the canonic process $\{B_t : t \geq 0\}$ is called G -Brownian motion.

The completion of $Lip(\Omega)$ under the Banach norm $\hat{\mathbb{E}}[\|\cdot\|]$ is denoted by $L_G^1(\Omega)$. $\hat{\mathbb{E}}[\cdot]$ can be extended uniquely to a sublinear expectation on $L_G^1(\Omega)$ (see Peng [12, 13, 14]).

We denote by $\mathcal{B}(\Omega)$ the Borel σ -algebra of Ω . It was proved in Hu and Peng [5] (see also Denis, Hu and Peng [3]) that there exists a weakly compact family $\mathcal{P}$ of probability measures defined on $(\Omega, \mathcal{B}(\Omega))$ such that

$$\hat{\mathbb{E}}[X] = \sup_{P \in \mathcal{P}} E_P[X], \quad \forall X \in L_G^1(\Omega).$$

For such weakly compact family $\mathcal{P}$, we can introduce the natural Choquet capacity

$$v(A) := \sup_{P \in \mathcal{P}} P(A), \quad A \in \mathcal{B}(\Omega).$$

Definition 2.4 A set $A \subset \Omega$ is polar if $v(A) = 0$. A property holds quasi-surely (q.s.) if it holds outside a polar set.

Definition 2.5 A real function X on Ω is said to be quasi-continuous if for each $\varepsilon > 0$, there exists an open set O with $v(O) < \varepsilon$ such that $X|_{O^c}$ is continuous.

The following proposition can be found in Denis, Hu and Peng [3].

Proposition 2.6 For each $X \in L_G^1(\Omega)$, there exists Y such that $Y = X$ q.s. and Y is quasi-continuous.

3 Main Theorem

In this section, we consider strict comparison theorem for Peng's G -expectation $\hat{\mathbb{E}}$, where $\hat{\mathbb{E}}$ defined on $(\Omega, L_G^1(\Omega))$ and there exists a weakly compact family $\mathcal{P}$ such that $\hat{\mathbb{E}}[X] = \sup_{P \in \mathcal{P}} E_P[X]$.

Theorem 3.1 *Let $X, Y \in L_G^1(\Omega)$ and $X \leq Y$ q.s. If*

$$\inf_{P \in \mathcal{P}} P(X < Y) > 0,$$

then $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$.

Proof. Since $\hat{\mathbb{E}}[X] - \hat{\mathbb{E}}[Y] \leq \hat{\mathbb{E}}[X - Y]$, we only consider the case of $X \leq 0$ q.s. and $\inf_{P \in \mathcal{P}} P(X < 0) > 0$.

For such X , we can choose ε such that $0 < \varepsilon < \inf_{P \in \mathcal{P}} P(X < 0)$. Since $X \in L_G^1(\Omega)$, by Proposition 2.6, there exists Y such that $X = Y$ q.s. and Y is quasi-continuous. Noting that $X = Y$ q.s. implies $\hat{\mathbb{E}}[X] = \hat{\mathbb{E}}[Y]$, and if $\hat{\mathbb{E}}[Y] < 0$ we also have $\hat{\mathbb{E}}[X] < 0$. Without loss of generality we can assume that X is quasi-continuous. By Definition 2.5, there exists an open set O such that $v(O) < \inf_{P \in \mathcal{P}} P(X < 0) - \varepsilon$ and $X|_{O^c}$ is continuous.

Set $A = \{\omega : X(\omega) \geq 0\}$, $A_n = \{\omega : X(\omega) \geq -1/n\}$, $F = A \cap O^c$ and $F_n = A_n \cap O^c$, then we can check that F_n is closed and $F_n \downarrow F$. Since $\mathcal{P}$ is weakly compact, we have $v(F_n) \downarrow v(F)$ (see Huber and Strassen [6]). There exists $n_0 \in \mathbb{N}$ such that $v(F_{n_0}) \leq v(F) + \varepsilon$.

We have

$$\begin{aligned} v(X \geq -1/n_0) - 1 &= v((A_{n_0} \cap O^c) \cup (A_{n_0} \cap O)) - 1 \\ &\leq v(F_{n_0}) + v(A_{n_0} \cap O) - 1 \\ &\leq v(F) + \varepsilon + v(O) - 1 \\ &\leq v(A) + \varepsilon + v(O) - 1 \\ &= -(\inf_{P \in \mathcal{P}} P(X < 0) - v(O) - \varepsilon) < 0 \end{aligned}$$

Since $X \leq 0$ q.s., we have $v(X > t) = 0$ for $t \geq 0$. Finally, we get

$$\begin{aligned} \hat{\mathbb{E}}[X] &= \sup_{P \in \mathcal{P}} E_P[X] = \sup_{P \in \mathcal{P}} \left(\int_0^\infty P(X > t) \mathfrak{t} + \int_{-\infty}^0 (P(X > t) - 1) \mathfrak{t} \right) \\ &\leq \int_0^\infty \sup_{P \in \mathcal{P}} P(X > t) \mathfrak{t} + \int_{-\infty}^0 (\sup_{P \in \mathcal{P}} P(X > t) - 1) \mathfrak{t} \\ &\leq \int_0^\infty v(X > t) \mathfrak{t} + \int_{-\infty}^0 (v(X \geq t) - 1) \mathfrak{t} = \int_{-\infty}^0 (v(X \geq t) - 1) \mathfrak{t} \\ &\leq \int_{-1/n_0}^0 (v(X \geq t) - 1) \mathfrak{t} \leq (v(X \geq -1/n_0) - 1)/n_0 < 0. \end{aligned}$$

□

In fact, the condition $\inf_{P \in \mathcal{P}} P(X < Y) > 0$ is not easy to verify. But for $X, Y \in Lip(\Omega)$, we can represent them by $X = \varphi(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$ and $Y = \psi(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$, where $\varphi, \psi \in C_{l,Lip}(\mathbb{R}^n)$ and B is the G -Brownian motion on $(\Omega, L_G^1(\Omega), \hat{\mathbb{E}})$. When $\underline{\sigma} > 0$, we have the following results.

Lemma 3.2 Suppose $\underline{\sigma} > 0$. Let $X \in Lip(\Omega)$ with the form $X = \varphi(B_1)$ and $\varphi(x) \leq 0, \forall x \in \mathbb{R}$. If there exists $x_0 \in \mathbb{R}$ such that $\varphi(x_0) < 0$, then $\hat{\mathbb{E}}[X] < 0$.

The proof of this lemma depends on some deep estimates of fully nonlinear parabolic partial different equations, which initially obtained by Krylov and Safonov [8], we will prove it in Appendix.

Theorem 3.3 Let $\underline{\sigma} > 0$ and $X, Y \in Lip(\Omega)$ with the forms $X = \varphi(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$ and $Y = \psi(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$, where $\varphi(x) \leq \psi(x), \forall x \in \mathbb{R}^n$. Then $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$ if and only if there exists $x_0 \in \mathbb{R}^n$ such that $\varphi(x_0) < \psi(x_0)$.

Proof. The necessity is obviously, we only need to prove the sufficiency.

We first consider the case of $X = \varphi(B_{t_1}, B_{t_2} - B_{t_1})$ and $Y = \psi(B_{t_1}, B_{t_2} - B_{t_1})$ with $\varphi \leq \psi$ and there exists $(x_1, x_2) \in \mathbb{R}^2$ such that $\varphi(x_1, x_2) < \psi(x_1, x_2)$. By Lemma 3.2, we have $\hat{\mathbb{E}}[\varphi(x, B_{t_2} - B_{t_1})] \leq \hat{\mathbb{E}}[\psi(x, B_{t_2} - B_{t_1})]$ for each $x \in \mathbb{R}$ and $\hat{\mathbb{E}}[\varphi(x_1, B_{t_2} - B_{t_1})] < \hat{\mathbb{E}}[\psi(x_1, B_{t_2} - B_{t_1})]$. Let us use Lemma 3.2 again, we conclude $\hat{\mathbb{E}}[\varphi(B_{t_1}, B_{t_2} - B_{t_1})] < \hat{\mathbb{E}}[\psi(B_{t_1}, B_{t_2} - B_{t_1})]$.

For $X = \varphi(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$ and $Y = \psi(B_{t_1}, B_{t_2} - B_{t_1}, \dots, B_{t_n} - B_{t_{n-1}})$, we repeat above procedure, thus $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$. $\square$

The condition of $\underline{\sigma} > 0$ is necessary. When $\bar{\sigma} = 0$, the above strict comparison theorem does not hold.

Example 3.4 Let $\underline{\sigma} = 0$ and $X = B_1 \wedge 0$, then we can check that $u(t, x) = x \wedge 0$ is the unique viscosity solution of G -heat equation: $\partial_t u = \bar{\sigma}^2 (\partial_{xx} u)^+ / 2$ with initial condition $u(0, x) = x \wedge 0$. But we have $\hat{\mathbb{E}}[X] = u(1, 0) = 0$. The strict comparison theorem does not hold.

Corollary 3.5 Let B_t be a G -Brownian motion with $\bar{\sigma}^2 = \hat{\mathbb{E}}[B_1^2] \geq -\hat{\mathbb{E}}[-B_1^2] = \underline{\sigma}^2 > 0$. Then we have

$$\inf_{P \in \mathcal{P}} P(a \leq B_t \leq b) > 0,$$

where $-\infty \leq a < b \leq \infty$ and $t > 0$.

Remark 3.6 Suppose $\underline{\sigma} > 0$ and $X, Y \in Lip(\Omega)$ satisfy the same conditions in Theorem 3.3. By Corollary 3.5, we can prove that $\inf_{P \in \mathcal{P}} P(X < Y) > 0$, so by Theorem 3.1, we also have $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$.

In the end, we give another form of strict comparison theorem.

Theorem 3.7 Let $X, Y \in L_G^1(\Omega)$ and $X \leq Y$ q.s.. If $v(X < Y) > 0$ and X has mean certainty, i.e., $\hat{\mathbb{E}}[X] = -\hat{\mathbb{E}}[-X]$, then $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$.

Proof. Since $X \leq Y$ q.s. and $v(X < Y) > 0$, there exists $P \in \mathcal{P}$ such that $P(X \leq Y) = 1$ and $P(X < Y) > 0$. Therefore $E_P[X] < E_P[Y]$. Since $\hat{\mathbb{E}}[X] = -\hat{\mathbb{E}}[-X]$, we have

$$\hat{\mathbb{E}}[X] = E_P[X] < E_P[Y] \leq \hat{\mathbb{E}}[Y].$$

□

But unfortunately, in general, if $X \leq Y$ and $v(X < Y) > 0$, it does not imply $\hat{\mathbb{E}}[X] < \hat{\mathbb{E}}[Y]$. We give the following counterexample, in which we will use the notion of quadratic variation process of G -Brownian motion $\langle B \rangle_t$. More details about this process can be found in Peng [10, 11, 12, 13, 14].

Example 3.8 Let B_t be a G -Brownian motion with $\bar{\sigma}^2 = \hat{\mathbb{E}}[B_1^2] > \underline{\sigma}^2 = -\hat{\mathbb{E}}[-B_1^2] > 0$. we consider $\langle B \rangle_t$ and $\bar{\sigma}^2 t$, then we have $\langle B \rangle_t \leq \bar{\sigma}^2 t$ q.s., and we can choose $P \in \mathcal{P}$ such that B_t becomes the classical Brownian motion with $E_P[B_1^2] = \underline{\sigma}^2$. Under this P , the quadratic variation process $\langle B \rangle_t$ equals $\underline{\sigma}^2 t$ P -a.s., so we have $P(\langle B \rangle_t < \bar{\sigma}^2 t) = 1$. It is easy to get $v(\langle B \rangle_t < \bar{\sigma}^2 t) = 1$, but $\hat{\mathbb{E}}[\langle B \rangle_t] = \bar{\sigma}^2 t = \hat{\mathbb{E}}[\bar{\sigma}^2 t]$.

A Proof of Lemma 3.2

In this section, we complete the proof of Lemma 3.2. We always suppose $\underline{\sigma} > 0$.

The following lemma is a spacial case of Theorem 1.1 in Krylov and Safonov [8]. We denote $B_R(x^0) := \{x : |x - x^0| < R\}$, $B_R := B_R(0)$ and $Q_{\theta,R} := (0, \theta R^2) \times B_R$.

Lemma A.1 Let $\theta > 1$ and $R \leq 2$, $u \in C^{1,2}(Q_{\theta,R})$, $u \geq 0$ be such that

$$\partial_t u - a(t, x) \partial_{xx} u = 0, \quad \text{in } Q_{\theta,R},$$

where $a \in L^\infty((0, \infty) \times \mathbb{R})$ and for some $\lambda > 0$, $\lambda^{-1} \leq a(t, x) \leq \lambda$, $\forall (t, x) \in (0, \infty) \times \mathbb{R}$.

Then there is a constant C depending only on λ, θ such that

$$u(\theta R^2, x) \geq C u(R^2, 0), \quad \forall x \in B_{R/2}.$$

We now give the proof of Lemma 3.2.

Proof of Lemma 3.2. We first consider the case of $\varphi \in C_{b,Lip}(\mathbb{R})$ (bounded and Lipschitzian continuous). Then the G -heat equation (1) has the unique classical solution, i.e., $u(t, x) \in C^{1,2}((0, 2) \times \mathbb{R})$ (see Krylov [7]). Since $\varphi(x) \leq 0$ for all $x \in \mathbb{R}$, by the well-known maximal principle, we have $u(t, x) \leq 0$. Since $\varphi(x_0) < 0$, by the continuity of $u(t, x)$, there exists $\varepsilon > 0$ and $(t_\varepsilon, x_\varepsilon) \in (0, 1/2) \times \mathbb{R}$, such that $u(t_\varepsilon, x_\varepsilon) < -\varepsilon$.

We set $v(t, x) = -u(t, x_\varepsilon - Mx)$, where $M > 2|x_\varepsilon|/\sqrt{t_\varepsilon}$. It is easy to verify that $v(t, x)$ is the unique solution of the following PDE:

$$\begin{cases} \partial_t v - a(t, x) \partial_{xx} v = 0, & (t, x) \in (0, 2) \times \mathbb{R}, \\ v|_{t=0} = -\varphi(x_\varepsilon - Mx), & x \in \mathbb{R}, \end{cases}$$

where

$$a(t, x) = \begin{cases} \bar{\sigma}^2/2M^2 & \text{for } (t, x) \text{ such that } \partial_{xx} u(t, x_\varepsilon - Mx) \geq 0, \\ \underline{\sigma}^2/2M^2 & \text{otherwise.} \end{cases}$$

Then $v(t, x)$ satisfies all the condition of above Lemma. We get $v(1, x_\varepsilon/M) \geq Cv(t_\varepsilon, 0) = -Cu(t_\varepsilon, x_\varepsilon) > 0$, where $C > 0$ depending on $\bar{\sigma}, \underline{\sigma}, x_\varepsilon, t_\varepsilon$. So we have $\hat{\mathbb{E}}[\varphi(B_1)] = u(1, 0) = -v(1, x_\varepsilon/M) < 0$.

For each $\varphi \in C_{l,Lip}(\mathbb{R})$ with $\varphi(x) \leq 0$ and $\varphi(x_0) < 0$, we can choose $\varphi' \in C_{b,Lip}(\mathbb{R})$ such that $\varphi(x) \leq \varphi'(x) \leq 0$ and $\varphi'(x_0) = \varphi(x_0) < 0$, then we have

$$\hat{\mathbb{E}}[\varphi(B_1)] \leq \hat{\mathbb{E}}[\varphi'(B_1)] < 0.$$

□

Acknowledgements

The author would like to thank Professor S. Peng for his helpful discussions.

References

- [1] G. Choquet, Theory of capacities. Annals det Institut Fourier 5 (1953) 131-195.
- [2] M. Crandall, H. Ishii., P.L. Lions, User's guide to viscosity solutions of second order partial differential equations. Bulletin of the American Mathematical Society 27 (1) (1992), 1-67.
- [3] L. Denis, M. Hu., S. Peng, Function spaces and capacity related to a sublinear expectation: application to G-Brownian Motion Pathes, see arXiv:0802.1240v1 [math.PR] 9 Feb 2008.
- [4] L. Gu, Second Order Parabolic Partial Differential Equations, Xiamen university press, Xiamen (Chinese version)(2002).
- [5] M. Hu, S. Peng, On representation theorem of G-expectations and paths of G-Brownian motion. Acta Mathematicae Applicatae Sinica 25 (3) (2009) 539-546.
- [6] P. Huber, V. Strassen, Minimax tests and the Neyman-Pearson Lemma for capacity. The Annals of Statistics, Vol. 1, No. 2 (1973) 252-263.
- [7] N.V. Krylov, Nonlinear Parabolic and Elliptic Equations of the Second Order, Reidel Publishing Company (Original Russian version by Nauka, Moscow, 1985).
- [8] N.V. Krylov, M.V. Safonov, A Certain property of solutions of parabolic equations with measurable coefficients. Izvestia Akad. Nauk. SSSR 40. (1980)
- [9] S. Peng, Backward SDE and related g-expectations, in Backward Stochastic Differential Equations, Pitman Research Notes in Math. Series, No.364, El Karoui Mazliak edit (1997) 141-159.

- [10] S. Peng, G -Expectation, G -Brownian Motion and Related Stochastic Calculus of Itô's type, preprint (pdf-file available in arXiv:math.PR/0601035v1 3Jan 2006), to appear in *Proceedings of the 2005 Abel Symposium*.
- [11] S. Peng, G -Brownian Motion and Dynamic Risk Measure under Volatility Uncertainty, Preprint: arXiv:0711.2834v1 [math.PR] 19 Nov 2007.
- [12] S. Peng, Multi-Dimensional G -Brownian Motion and Related Stochastic Calculus under G -Expectation, in Stochastic Processes and their Applications, **118**(12) (2008) 2223-2253.
- [13] S. Peng, Survey on normal distributions, central limit theorem, Brownian motion and the related stochastic calculus under sublinear expectations, Science in China Series A: Mathematics, **52**, No.7 (2009) 1391–1411.
- [14] S. Peng, Nonlinear Expectations and Stochastic Calculus under Uncertainty –with Robust Central Limit Theorem and G -Brownian Motion, arXiv:1002.4546v1 [math.PR] 24 Feb 2010.